

Meet Michael

Learning Outcomes

- Individualising glycaemic treatment targets depending on patient wishes, comorbidities and hypoglycaemia risk
- Determine when to escalate to triple oral therapy and insulin treatments
- Diagnosis and management of diabetes-related complications, particularly cardiovascular risk
- Implications of insulin therapy for commercial drivers



VISIT ONE

Michael is a 58-year old male. He was diagnosed with type 2 diabetes six years ago with a background history of diastolic hypertension for ten years. He was unable to tolerate metformin or metformin XR due to severe gastrointestinal side effects. A sulfonylurea was commenced five years ago and a DPP-4 inhibitor added two years ago. He was also prescribed an ACE inhibitor for hypertension and mild proteinuria five years ago. Screening for complications is up-to-date with no evidence of peripheral neuropathy or retinopathy.

Michael works as a driver for a long haul trucking company. His job is physically active as he loads and unloads the truck with each delivery. He states that he would prefer to avoid insulin therapy unless absolutely necessary. He smoked a pack of cigarettes a day for the 30 years but ceased five years ago. He drinks two-four standard drinks of alcohol on weekends. He is adopted and unaware of his biological family's medical history.

What are the management issues for this patient?

- Patient's age and medical co-morbidities suggest that an HbA_{1c} target of 53 mmol/mol (7%) would be appropriate
- Individualising glycaemic treatment targets depending on patient wishes, comorbidities and hypoglycaemia risk
- Selection of triple therapy to improve glycaemic control
- Assessment of absolute cardiovascular risk
- Optimal management of high blood pressure

Current medications

Glimepiride 4mg daily
Saxagliptin 5mg daily
Perindopril 4mg daily

Allergies

Nil known drug allergies

Examination

Blood pressure 145/85 mmHg
Weight 87 kg, Height 180 cm, BMI 27 kg/m²
Ankle jerks present, monofilament sensation intact, pedal pulses present, nil evidence of ulceration

Investigations

HbA_{1c} 69 mmol/mol (8.5%)
eGFR 55 ml/min/m²

What is your management plan?

1. Patient's age and lack of medical co-morbidities suggest that an HbA_{1c} target of 53 mmol/mol (7%) would be appropriate
2. Stop glimepiride. Start gliclazide MR 120 mg; gliclazide has a lower risk of hypoglycemia than glimepiride given differences in active metabolites
3. Stop DPP-4 inhibitor. Start empagliflozin. DPP-4 inhibitors and SGLT2 inhibitors are not PBS-reimbursed for use together
4. Ensure patient is educated regarding blood glucose testing before driving
5. Ensure the road traffic authority is informed of Michael's diabetes
6. Measure lipid profile and urine albumin/creatinine ratio (ACR)

VISIT TWO

When Michael is reviewed in two weeks, he has improved his diet. Michael's friend Alec has recently had a triple coronary bypass, and although he has not had any symptoms, Michael is concerned about his own risk of heart disease.

Current medications

Gliclazide MR 120 mg daily
Empagliflozin 25mg daily
Perindopril 4mg daily

What are the management issues for this patient?

- Calculation of absolute cardiovascular risk
- Prevention of cardiovascular disease

Examination

Blood pressure 138/80 mmHg
Weight 87 kg

Investigations

ECG – Sinus 65 beats per minute, nil abnormalities detected
Total Cholesterol 4.5 mmol/L, HDL 1.2 mmol/L, LDL 3.3 mmol/L
TG 1.5 mmol/L
Urine ACR 5.5 mg/mmol

The Australian absolute cardiovascular risk calculator is used to estimate Michael's cardiovascular risk and indicates a 12% risk in the next five years.

What is your management plan?

1. Provide information on prevention of cardiovascular disease.
2. Commence statin therapy.
3. Monitor blood pressure and titrate anti-hypertensive therapy.

VISIT THREE

Michael is reviewed in three months, HbA_{1c} continues to be above target. His home blood glucose recordings indicate some elevation in fasting and pre-prandial glucose levels (7.5-9.0 mmol/L). What is the next line of therapy?

Current medications

Gliclazide MR 120mg daily
Empagliflozin 25mg daily

What are the management issues for this patient?

- Intensification of diabetes therapies
- Commencement of insulin, factors to consider in determining the optimal insulin regimen for each patient taking in account convenience, flexibility, medication adherence, lifestyle and patient preference

What is your management plan?

1. Commence basal insulin as HbA_{1c} target has not been achieved with the dual oral therapy.

Perindopril 4mg daily
Atorvastatin 40mg daily

Examination

Blood pressure 120/80 mmHg
Weight 83 kg, Height 180 cm, BMI 25.6 kg/m²

Investigations

HbA_{1c} 62 mmol/mol (7.8%)

2. Sulfonylurea and SGLT2 inhibitor may be continued to reduce insulin requirements.
3. Referral to diabetes nurse educator for information on insulin injecting technique, hypoglycaemia management and prevention while working and insulin storage on the road. Dietitian review to ensure that Michael is evenly distributing his carbohydrate intake to reduce the variability in his BGLs.
4. Ensure the driving authority has been informed of insulin therapy commencement.

VISIT FOUR

Three months later, Michael presents for follow up. He has seen an endocrinologist for review of his diabetes. He is tolerating his medications and reports no episodes of hypoglycaemia. His ability

to hold a commercial driving license is not affected as his control has improved on the new regimen with an HbA_{1c} of 55mmol/mol (7.2%).

Additional resources

http://www.cvdcheck.org.au/pdf/Absolute_CVD_Risk_Full_Guidelines.pdf

<http://www.cvdcheck.org.au>

<https://www.diabetesaustralia.com.au/driving>

Australian absolute cardiovascular disease risk calculator

Enter patient information below:

Sex Male Female
Age 58 years
Systolic blood pressure 138 mmHg
Smoking status Yes No
Total cholesterol 4.5 mmol/L
HDL cholesterol 1.2 mmol/L
Diabetes Yes No
ECG LVH Yes No Unknown

Your heart and stroke risk score is **12%**

This means you are at moderate (medium) risk of getting cardiovascular disease in the next 5 years.

Click here if you would like to have a look at the information on this website that explains what your risk score means.

The next step is to talk to your doctor about what steps you can take to lower your chance of getting cardiovascular disease.

Please note: the absolute risk calculator score is only a guide to your heart and stroke risk score. Print out this page and take it to your doctor for further information on your personal risk.

View guidelines and resources

Diabetes Australia
KIDNEY HEALTH AUSTRALIA
Heart Foundation
strokefoundation

An initiative of the National Vascular Disease Prevention Alliance