

References:

1. UKPDS Group. Lancet 1998;352:854-65.
2. UKPDS Group. Lancet 1998;352:837-53.
3. ADVANCE Collaborative Group. NEJM 2008;358:2560-72.
4. White WB, et al. NEJM 2013;369:1327-35.
5. Zannad F, et al. Lancet 2015;385:2067-76.
6. Scirica BM, et al. NEJM 2013;369:1317-26.
7. Scirica BM, et al. Circulation 2014;130:1579-88.
8. Green JB, Bethel MA, et al. NEJM 2015;373:232-42.
9. Meier JJ, et al. Diabetologia 2014;57:1320-1324.
10. McGill JB, et al. Diabetes Care 2013;36:237-44.
11. Dormandy JA, et al. Lancet 2005;366:1279-89.
12. Home PD, et al. Lancet 2009;373 :2125-35.
13. Zinman B, et al. NEJM 2015;372:2117-28.
14. Pfeff MA, et al. Symposium, 75th Scientific Sessions of the American Diabetes Association; Boston, MA; 2015.
15. Bentley-Lewis R, et al. Am Heart J 2015;169:631-38.
16. Marso SP, et al. NEJM 2016;375:311-322.
17. ORIGIN Trail Investigators. NEJM 2012; 367:319-328.